Abbreviazioni

SV stato vegetativo

SMC stato di minima coscienza

E-SMC emergenza dallo stato di minima coscienza

Disorders of Consciousness DOC

P-DOC **Prolonged-Disorders of Consciousness**

MATADOC Music Therapy Assessment Tool for Awareness in Disorders of Consciousness

Bibliografia

1. Alberto Ezzu, Roberto Messaglia

Introduzione alla musicoterapia: Storia, fondamenti, modelli, applicazioni cliniche, glossario.

Ed. Musica Pratica 2006

Collana di Musicoterapia a cura del Centro Musicoterapia Benenzon Italia

2. AAVV

Musica tra neuroscienze arte e terapia

Ed. Musica Pratica 2007

Collana di Musicoterapia a cura del Centro Musicoterapia Benenzon Italia

3. Shantala Hegde

Music-based cognitive remediation therapy for patients with traumatic brain injury. Frontiers in neurology - MINI REVIEW ARTICLE published: 24 March 2014

- 4. Hurt CP, Rice RR, McIntosh GC, ThautMH. Rhythmic auditory stimulation in gait training for patients with traumatic brain injury. J MusicTher (1998) 35: 228–41.
- 5. Music therapy for acquired brain injury (Review) Bradt J, Magee WL, Dileo C, Wheeler BL, McGilloway E The Cochrane Library 2010, Issue 7.
- 6. Thaut MH, Gardiner JC, Holmberg D, Horowitz J, Kent L, Andrews G, et al. Neurologic music therapy improves executive function and emotional adjustment in traumatic brain injury rehabilitation.

Ann N Y Acad Sci (2009) **1169**:406–16.

- 7. Särkämö T, Tervaniemi M, Laitinen S, Forsblom A, Soinila S, Mikkonen M, et al. Music listening enhances cognitive recovery and mood after middle cerebral artery stroke. Brain (2008) 131:866-76.
- 8. Royal College of Physicians. 2013.

Prolonged Disorders of

Consciousness: National Clinical Guidelines.

London: RCP.

9. Wendy L. Mageel and Julian O'Kelly

Music therapy with disorders of consciousness: current evidence and emergent evidencebased practice.

Ann. N.Y. Acad. Sci. 1337 (2015) 256-262

10. Schnakers, C., et al.

Diagnostic accuracy of the vegetative and minimally conscious state: clinical consensus versus

standardized neurobehavioral assessment.

BMC Neurol. (2009) 9:35

11. Laureys, S., A.M. Owen & N.D. Schiff.

Brain function in coma, vegetative state, and related disorders.

Lancet Neurol. (2004) 3: 537-546.

12. Laureys, S., et al.

Restoration of thalamocortical connectivity after recovery from persistent vegetative state. *Lancet* (2000).**355**: 1790–1791.

13. Whyte, J., *et al*.

Predictors of outcome in prolonged posttraumatic disorders of consciousness and assessment of

medication effects: a multicenter study.

Arch. Phys. Med.Rehabil. 2005.86: 453–462.

14. Laureys, S., et al.

Auditory processing in the vegetative state.

Brain 2000. 123(Pt 8): 1589-1601.

15. Schiff, N.D., et al.

Residual cerebral activity and behavioural fragments can remain in the persistently vegetative

brain.

Brain 2002.125: 1210-1234.

16. Kassubek, J., et al.

Activation of a residual cortical network during painful stimulation in long-term postanoxic vegetative state: a 15O-H2O PET study.

J. Neurol. Sci. 2003.212: 85-91.

17. Kotchoubey, B.

Apallic syndrome is not apallic: is vegetative state vegetative?

Neuropsychol. Rehab. . 2005. 15: 333-356.

18. Coleman, M.R., et al.

Do vegetative patients retain aspects of language comprehension? Evidence from fMRI. *Brain* 2007. **130**: 2494–2507.

19. Monti, M., et al.

Wilful modulation of brain activity in disorders of consciousness.

N. Engl. J.Med. 2010.362: 579–589.

20. John, E.R., et al.

Source imaging of QEEG as a method to detect awareness in a person in vegetative state. *Brain Injury* 2011.**25**: 426–432.

21. Owen, A.M., et al.

Using a hierarchical approach to investigate residual auditory cognition in persistent vegetative

state.

Prog. Brain Res. 2005.**150**: 457-471.

22. Owen, A.M., et al.

Detecting awareness in the vegetative state.

Science 2006. 313: 1402.

23. Boly, M., et al.

Cerebral processing of auditory and noxious stimuli in severely brain injured patients: differences between VS and MCS.

Neuropsychol. Rehabil. 2005.15: 283-289.

24. Perrin, F., *et al*.

Brain response to one's own name in vegetative state, minimally conscious state, and locked-in

syndrome.

Arch. Neurol. 2006.63: 562-569.

25. Jones, S.J., et al.

Auditory evoked potentials to spectrotemporal modulation of complex tones in normal subjects

and patients with severe brain injury.

Brain 2000.123: 1007-1016.

26. Gill-Thwaites, H. & R.Munday. 2004.

The sensorymodality assessment and rehabilitation technique (SMART): a valid and reliable assessment for vegetative state and minimally conscious state patients.

Brain Injury **18**: 1255–1269.

27. Laureys, S., *et al*.

Cerebral processing in the minimally conscious state.

Neurology 2004. 63: 916–918.

28. Laureys, S., F. Perrin & S. Bredart.

Self-consciousness in non-communicative patients.

Conscious. Cogn. 2007. 16: 722–741; discussion 742–725.

29. Schnakers, C., et al.

Detecting consciousness in a total locked-in syndrome: an active event-related paradigm. *Neurocase* 2009. **15**: 271–277.

30. O'Kelly, J., *et al*.

Neurophysiological and behavioral responses to music therapy in vegetative and minimally conscious states.

Front. Hum. Neurosci. 2013.7: 884.

31. S"ark"am"o, T., et al.

Music listening enhances cognitive recovery and mood aftermiddle cerebral artery stroke. *Brain* 2008.**131**: 866–876.

32. S"ark"am"o, T., et al.

Structural changes induced by daily music listening in the recovering brain after middle cerebral

artery stroke: a voxel-based morphometry study.

Front. Hum. Neurosci. 2014. 8: 245.

33. Magee, W.L., et al.

Music Therapy Assessment Tool for Awareness in Disorders of Consciousness (MATADOC):

standardisation of the principal subscale to assess awareness in patients with disorders of consciousness.

Neuropsychol. Rehabil. 2014. 24: 101–124.

34. Shiel, A. & B.A. Wilson.

Can behaviours observed in the early stages of recovery after traumatic brain injury predict poor outcome?

Neuropsychol. Rehabil. 2005. 15: 494-502.

35. Vanhaudenhuyse, A., et al.

Assessment of visual pursuit in post-comatose states: use a mirror.

J. Neurol. Neurosurg. Psychiatry 2008.79: 223.

36. Seel, R.T., et al.

Assessment scales for disorders of consciousness: evidence-based recommendations for clinical

practice and research.

Arch. Phys.Med. Rehabil. 2010.91: 1795-1813.

37. Giacino, J.T., K. Kalmar & J. Whyte.

The JFKComa Recovery Scale-Revised: measurement characteristics and diagnostic utility.

Arch. Phys. Med. Rehabil. 2004.85: 2020–2029.

38. Gill-Thwaites, H. & R. Munday. 2008.

SMART: Sensory Modality Assessment and Rehabilitation Technique Manual.

London: Royal Hospital for Neuro-disability.

39. Shiel, A., et al.

Thewessex head injurymatrix (WHIM) main scale: a preliminary report on a scale to assess andmonitor patient recovery after severe head injury.

Clin. Rehabil. 2000. 14: 408-416.

40. Rappaport, M., A.M. Dougherty & D.L. Kelting.

Evaluation of coma and vegetative states.

Arch. Phys.Med. Rehabil. 1992.73: 628-634.

41. O'Kelly, J. & W.L. Magee.

The complementary role of music therapy in the detection of awareness in disorders of consciousness: an audit of concurrent SMART and MATADOC assessments. *Neuropsychol. Rehabil.* 2013. **23**: 287–298.

42. Cicerone KD, Dahlberg C, Malec JF, Langenbahn DM, Felicetti T, Kneipp S, et al. Evidence-based cognitive rehabilitation: updated review of the literature from 1998 through 2002.

Arch Phys Med Rehabil (2005) **86**:1681–92.

- 43. Cicerone KD, Langenbahn DM, Braden C, Malec JF, Kalmar K, Fraas M, et al. Evidence-based cognitive rehabilitation: updated review of the literature from 2003 through 2008. *Arch Phys Med Rehabil* (2011) **92**:519–30.
- 44. Rohling ML, Faust ME, Beverly B, Demakis G. Effectiveness of cognitive rehabilitation following acquired brain injury: a meta-analytic reexamination of Cicerone et al.'s (2000,2005) systematic reviews. *Neuropsychology* (2009) **23**:20.
- 45. Cappa SF, Benke T, Clarke S, Rossi B, Stemmer B, Van Heugten CM. EFNS guidelines on cognitive rehabilitation: report of an EFNS task force. *Eur J Neurol* (2005) **12**:665–80.
- 46. Cernich AN, Kurtz SM, Mordecai KL, Ryan PB. Cognitive rehabilitation in traumatic brain injury. *Curr Treat Option sNeurol* (2010) **12**:412–23.
- 47. van Heugten C, Gregorio GW, Wade D.

Evidence-based cognitive rehabilitation after acquired brain injury: a systematic review of content of treatment.

Neuropsychol Rehabil (2012) **22**:653–73.

48. Thaut MH.

Neurologic music therapy in cognitive rehabilitation.

Music Percept (2010) **27**:281–5.

49. Schlaug G.

PartVI introduction: listening to and making music facilitates brain recovery processes. *AnnNYAcadSci* (2009) **1169**:372–3.

50. Peretz I, Zatorre RJ.

Brain organization for music processing. *Annu Rev Psychol* (2005) **56**:89–114.

51. Zatorre R, McGill J.

Music, the food of neuroscience?

Nature (2005) **434**:312–5.

52. Levitin DJ, Tirovolas AK.

Current advances in the cognitive neuroscience of music.

Ann N Y Acad Sci (2009) 1156:211-31.

53. Patel AD.

Music, biological evolution, and the brain.

In: Bailar M, editor. Emerging Disciplines.

Houston, TX: Rice University Press (2010).p.91–144.

54. Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ.

Anatomically distinct dopamine release during anticipation and experience of peak emotion to music.

Nat Neurosci (2011) 14:257-62.

55. Blood AJ, Zatorre RJ.

Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion.

Proc Natl Acad Sci U S A (2001) 98:11818-23.

56. Menon V, Levitin DJ.

The rewards of music listening :response and physiological connectivity of the mesolimbic system.

Neuroimage (2005) 28:175-84.

57. Salimpoor VN, vanden Bosch I, Kovacevic N, McIntosh AR, Dagher A, Zatorre RJ. Interactions between the nucleus accumbens and auditory cortices predict music reward value. *Science* (2013) **340**:216–9.

Frontiers in Neurology | Neurotrauma March2014|Volume5|Article34| 6

58. Koelsch S, Fuermetz J, Sack U, Bauer K, Hohenadel M, Wiegel M, et al.

Effects of music listening on cortisol levels and propofol consumption during spinal anaesthesia.

Front Psychol (2011) 2:58.

59. Suda M, Morimoto K, Obata A, Koizumi H, Maki A.

Emotional responses to music: towards scientific perspectives on music therapy. *Neuroreport* (2008) **19**(1):75–8.

60. Khalfa S, Bella SD, Roy M, Peretz I, Lupein SJ.

Effects of relaxing music on salivary cortisol level after psychological stress. *AnnNYAcadSci* (2003) **999**:374–6.

61. Chanda ML, Levitin DJ.

The neurochemistry of music.

TrendsCognSci (2013) 17:179–93.

62. Fukui H, Toyoshima K.

Music facilitate the neurogenesis, regeneration and repair of neuron.

MedHypotheses (2008) **71**:765–9.

63. Thaut MH.

Music in Therapy and Medicine: From Social Science to Neuroscience.

NewYork: Routledge; Taylor and Francisgroup(2005).

64. Jeong E, Lesiuk TL.

Development and preliminary evaluation of a music-based attention assessment for patients with traumatic brain injury.

J MusicTher (2011) 48:551–72.

65. Knox R, Yokota-Adachi H, Kreshner J, Jutai J.

Musical attention training program and alternating attention in brain injury: an initial report. *MusicTher Perspect* (2003) **21**:99–104.

66. Knox R, Jutai J.

Music-based rehabilitation of attention following brain injury. *Can JRehabil* (1996) **9**:169–81.

67. Bradt J, Magee WL, Dileo C, Wheeler BL, Mc Gilloway E.

Musictherapy for acquired brain injury.

Cochrane Database Syst Rev (2010) 7:CD006787.

68. del'Etoile SK.

Neurologic music therapy: a scientific paradigm for clinical practice. *MusicMed* (2010) **2**:78–84.

69. Clair AA, Pasiali V, Lagasse B.

Neurologic music therapy. 2nd ed.

In: Darrow AA, editor. Introduction to Approaches in Music Therapy.

SilverSpring, MD: The American Music Therapy Association (2008).p.153–72.

70. Thaut MH.

Neurologic music therapy in sensori motor rehabilitation.

In: Thaut MH, editor. Rhythm, Music and the Brain.

NewYork: Routledge; Taylorand Francis group (2005). p. 137-64.

71. Thaut MH.

Neurologic music therapy in speech and language rehabilitation.

In: Thaut MH, editor. Rhythm, Music and the Brain.

New York: Routledge; Taylor and Francis group (2005). p.165–78.

72. Thaut MH.

Neurologic music therapy in cognitive rehabilitation.

In: Thaut MH, editor. Rhythm, Music and the Brain.

New York: Routledge; Taylor and Francis group (2005). p.179–202.